



XLVI Congresso Nazionale AMCLI

11 - 14 Novembre 2017

Palacongressi di Rimini

Corso Precongressuale

Le infezioni da non dimenticare nel paziente trapiantato

Adenovirus

Piralla Antonio



Fondazione IRCCS
Policlinico San Matteo

Sistema Socio Sanitario



Regione
Lombardia

*Molecular Virology Unit, Microbiology and Virology Department
Fondazione IRCCS Policlinico San Matteo, Pavia*

Infections in Immunocompromised Hosts and Organ Transplant Recipients: Essentials

Jay A. Fishman

Transplant Center, Transplant Infectious Disease and Compromised Host Service, Infectious Disease Division, Massachusetts General Hospital, Harvard Medical School, Boston, MA

17:S34-S37, 2011

36 FISHMAN

LIVER TRANSPLANTATION, November 2011




Figure 1. Timeline of posttransplant infections.



Stem Cell
Research & Therapy


Kedia et al., J Stem Cell Res Ther 2013, S3
<http://dx.doi.org/10.4172/2157-7633.S3-002>

Review Article

Open Access

Infectious Complications of Hematopoietic Stem Cell Transplantation

Shiksha Kedia¹, Pranab Sharma Acharya¹, Farhan Mohammad¹, Huy Nguyen¹, Deepak Asti¹, Suchita Mehta^{1*}, Manisha Pant² and Neville Mobaraka³



HAdV infections in HSCT and SOT: State of art

SOT



- ✓ Adenovirus (HAdV) infections have been reported in wide range of SOT population, including those receiving heart, lung, liver intestinal and kidney transplants. (*Hoffman JA, Ped Transpl 2006 10:17-25*)
- ✓ Asymptomatic HAdV DNA-emia is common among SOT recipients and generally is not associated with progression to symptomatic disease. (*Humar et al. Am J Transplant 2005; 5:2555-2559*)

HSCT

- ✓ HAdV has the highest impact on morbidity, graft survival, and mortality in profoundly T cell-depleted transplant recipients, especially in children and in patients after HSCT (*Florescu DF, Hoffman JA. Am J Transplant 2013; 13(Suppl 4): 206–211.*)
- ✓ Poor and late reconstitution is a factor associated with increased risk of adenovirus disease in HSCTR. (*Moss and Rickinson Nature Reviews Immunology 2005; 5: 9–20*)


Viral structure and genome

- ✓ Linear, double-stranded DNA ($\approx 34\text{--}36$ kb in size)
- ✓ The virions are non-enveloped icosadeltahedrons
- ✓ The capsid comprises 240 capsomeres, which consist of hexons and pentons. The 12 pentons, which are located at each of the vertices, have a penton base and a fiber.
- ✓ There are currently over **60 HAdV types** in seven species (human adenovirus A–G), with HAdV-D containing the most members.




Antigenic sites

- ✓ Hexon, penton and fiber proteins are the most immunogenic parts of the HAdV capsid.
- ✓ Fiber and penton are targeted by neutralizing antibodies preventing viral entry, whereas antibodies directed to hexon prevent HAdV uncoating and genome delivery to the nucleus.
- ✓ Hexon is highly conserved across human HAdV species and has been used as an important antigen for monitoring of HAdV-specific T-cells.



HAdV pathogeneis

- ✓ Adenoviruses primarily infect children and less commonly infect adults.
- ✓ Spread by aerosol, close contact or oral-fecal route.
- ✓ Virus infects first mucoepithelial cells causing direct cell damage.
- ✓ Virus persist in lymphoid tissue
- ✓ Disease is determined by the tissue tropism of a HAdV type.



Clinical syndromes and HAdV types

- Acute respiratory disease (URTI and LRTI)
- Pharyngoconjunctival fever
- Epidemic keratoconjunctivitis
- Gastroenteritis and diarrhea

| Subgroup | Serotype | Sites of infection |
|----------|--|---------------------|
| A | 12, 18, 31 | Gastrointestinal |
| B1 | 3, 7, 16, 21, 50 | Respiratory |
| B2 | 11, 14, 34, 35 | Urinary tract/renal |
| C | 1, 2, 5, 6 | Respiratory |
| D | 8, 9, 10, 13, 15, 17, 19, 20, 22–30, 32, 33, 36, 37, 38, 39, 42–48, 49, 51 | Eye |
| E | 4 | Respiratory |
| F | 40, 41 | Gastrointestinal |

Journal of Clinical Virology 81 (2016) 78–81



Contents lists available at [ScienceDirect](#)

Journal of Clinical Virology

journal homepage: www.elsevier.com/locate/jcv



Case report

Acute respiratory distress syndrome in adenovirus type 4 pneumonia: A case report



R. Narra^a, P. Bono^a, A. Zoccoli^a, A. Orlandi^a, S. Piconi^b, G. Grasselli^c, S. Crotti^c, A. Girello^d,
A. Piralla^d, F. Baldanti^{d,e}, G. Lunghi^{a,*}

Clinical syndromes – Immunocompromised patients

Patients receiving stem cell transplantation and with T-cell deletion and lymphopenia are at risk for serious adenovirus infections.


Infections in these patients may cause variable manifestations including:

- ✓ hemorrhagic cystitis
- ✓ pneumonia
- ✓ nephritis
- ✓ gastroenteritis
- ✓ Disseminated infection with organ localization preceded by viremia

Infection occurs either by reactivation of old infection or by exogenous transmission from other infected hosts.

Laboratory diagnosis


- ✓ Samples types should be collected from a site or secretion relevant to the disease symptoms.
 - Respiratory samples (NPA, nasal swabs, BAL)
 - Conjunctival swabs or scrapings
 - urine
 - Stool
 - Peripheral Blood
 - Biopsy (organ localization)
- ✓ qReal-time PCR can be used to detect and quantify HAdV.
- ✓ Sequencing of Hexon gene for typing.
- ✓ Serologic testing is rarely used except for epidemiologic purposes.



HAdV-specific Immunity decrease with older age

- ✓ IgG level reflects the exposure of young people at HAdV infections.
- ✓ HAdV-specific T cells did not differ between transplant recipients and control subjects.
- ✓ The absolute frequencies and the percentage of individuals with detectable virus-specific T cells were significantly higher in younger individuals, compared with those of the older individuals.

T-CD4



IgG




Table 1. Adenovirus-specific T cell frequencies, by control subject or transplant patient status and age group.

| Group, T cell frequency | All individuals | Age group, years | | |
|---------------------------------|-----------------|------------------|-----------|------------------------|
| | | 18-39 | 40-60 | > 60 |
| Control subjects, <i>n</i> | 171 | 74 | 30 | 67 |
| > 0.05% | 53.2 (91) | 87.8 (65) | 53.3 (16) | 16.4 (11) ^a |
| ≤ 0.05% | 46.8 (80) | 12.2 (9) | 46.7 (14) | 83.6 (56) ^a |
| Transplant recipients, <i>n</i> | 59 | 16 | 27 | 16 |
| > 0.05% | 55.9 (33) | 87.5 (14) | 48.2 (13) | 37.5 (6) ^b |
| ≤ 0.05% | 44.1 (26) | 12.5 (2) | 51.9 (14) | 62.5 (10) ^b |

NOTE. Data are percentage (no.) of individuals with the indicated adenovirus-specific T cell frequency, unless otherwise indicated.

^a $P < .0001$, χ^2 test.


^b $P = .0094$, χ^2 test.

Sester et al., 2002 JID 185:1379-1387


HAdV-specific Immunity (Sester et al. Am J Transpl 2016;16 (6):1697-1706.C

Absolute lymphocyte counts $<300/\mu\text{L}$ predict poor outcome.

SOTR




Chemotherapy or HSCTR




Increasing lymphocyte counts and the appearance of ADV-specific CD4 and CD8 T cells have been correlated with clearance of viremia.

immunocompetent



HSCTR




HAdV-specific Immunity

- ✓ Detectable immunity reflects ongoing viral exposure, explaining the particular susceptibility of immunocompromised children for infectious complications.
- ✓ **The functional role of HAdV-specific T-cells** is supported by results of adoptive HAdV-specific T-cell transfer that successfully **prevented** or **cleared** HAdV replication in pediatric HSCT recipients.
(Papadopoulou A et al. *Sci Transl Med* 2014; 6(242): 242–283.)
- ✓ Little evidence supports surveillance for HAdV-specific T-cells after SOT, but its application would reasonably be focused on pediatric patients. (Florescu DF, et al. *Transplantation* 2010; 90(2): 198–204.)

Risk factor for HAdV disease

- ✓ Very young age (<5 years)
- ✓ T-cell depletion
- ✓ Immunosuppression

Treatment and Control




✓ Cidofovir (i.v) → ( side effects on kidney)

Risk of renal toxicity was 26% in HSCTR and most of toxicity was mild (Ljungman et al., Bone Marrow Transp 2003;31:481-486)

✓ Infusion of donor derived HAdV-specific T-Cells (only in experienced centers)


✓ Reduce immunosuppression (if possible)



Others

✓ Ribavirin (not recommended)

✓ Brancidofovir ?? Only case reports so far



Because of toxicity and limited effectiveness, cidofovir is still not ideal, but, when used as a **preemptive therapy**, it is a reasonably safe option to buy time until immune recovery.

HAdV DNA load: Frequency of monitoring

Early detection of HAdV viremia is helpful for starting therapy within a certain timeframe before the occurrence of disease symptoms.


The monitoring of HAdV load is based on stratification of patients based on risk.

- ✓ Low risk – HSCT - family haploidentical donor
No monitoring, only based on clinical signs
- ✓ Medium risk – HSCT – Match Unrelated Donor (MUD)
Ones at month up to to 100 days after TX
- ✓ High risk – HSCT- cord blood and T cell–depleted graft recipients
Ones a week up to 100 days after TX and

Treatment guideline: HSCT recipients


pre-emptive treatment

secondary treatment




Caroline A. Lindemans et al. Blood 2010;116:5476-5485

Adenovirus infection in HSCTR or in chemotherapy: Our experience 2015-2017




- ✓ A total of **35/172 (20.3%)** pediatric patients and **2/70 (2.9%)** adults patients had at least one positive blood sample.
- ✓ Among pediatric patients 33/35 (94.3%) had a single episode of HAdV infections, while 2/35 (5.3%) had two HAdV episodes.

In 14/37 (37.8%) HAdV episodes only one sample was positive, while in 23/37 (62.2%) episodes at least 2 samples were positive.



Adenovirus infection in HSCTR or in chemotherapy: Our experience 2015-2017

Median duration of HAdV episode: 20 days (range 3-90 days)




Adenovirus screening in pediatric HSCTR

Biol Blood Marrow Transplant 20 (2014) 250–256

Patient, Virus, and Treatment-Related Risk Factors in Pediatric Adenovirus Infection after Stem Cell Transplantation: Results of a Routine Monitoring Program


Martin Mynarek^{1,2}, Tina Ganzenmueller³, Annika Mueller-Heine⁴, Christopher Mielke¹, Andrea Gonnermann⁴, Rita Beier¹, Martin Sauer¹, Britta Eiz-Vesper^{2,5}, Ute Kohstall⁶, Karl-Walter Sykora¹, Albert Heim³, Britta Maecker-Kolhoff^{1,2,*}



- ✓ Single center 2003-2012 (GER)
- ✓ Prospective weekly HAdV screening (n=238)
 - Mostly type A31, C1, C2
- ✓ HAdV load >1000 DNA copies/ml in 15.5%
- ✓ Limited direct mortality 0.84%
- ✓ peak HAdV blood levels of >10,000 copies/mL were an independent risk factor of poor overall survival.
- ✓ Cidofovir i.v. treatment – Possibly stabilized/decreased HaDV load by 1 log₁₀

HAdV load with Cidofovir vs T-cell Recostitution

Biol Blood Marrow Transplant 21 (2015) 293–299



Biology of Blood and Marrow Transplantation

journal homepage: www.bbmt.org



The Effect of Cidofovir on Adenovirus Plasma DNA Levels in Stem Cell Transplantation Recipients without T Cell Reconstitution



Gertjan Lugthart*, Marloes A. Oomen, Cornelia M. Jol-van der Zijde, Lynne M. Ball, Dorine Bresters, Wouter J.W. Kollen, Frans J. Smiers, Clementien L. Vermont, Robbert G.M. Bredius, Marco W. Schilham, Maarten J.D. van Tol, Arjan C. Lankester

- ✓ Children (n=36 4,5 yrs) – pediatric allo HCT in Leiden (NL) 2003-2012
- ✓ Monitor weekly: HAdV DNA load, T-cells (CD3+, CD14-) and NK cells (CD56+, CD14-)
- ✓ HAdV clearance correlated with \uparrow T-cells
- ✓ Cidofovir at 1mg/kg 3x/wk only stabilizes HAdV load

Take home messages

- ✓ Children >>> Adult and HSCT >>> SOT
- ✓ T-cell reconstitution remains essential for viral clearance.
- ✓ Patients with HAdV viremia after HSCT might benefit from cidofovir treatment through a stabilization of the HAdV load but depending lymphocyte reconstitution for HAdV elimination.
- ✓ Therefore, adoptive T-cell therapy should be considered in patients not responding to cidofovir treatment.
- ✓ For clinical decision-making, the combined weekly monitoring of blood HAdV DNA levels and lymphocyte reconstitution provides an objective tool for the guidance of personalized antiviral treatment and prevention of unnecessary exposure to cidofovir.

Acknowledgements

Fondazione IRCCS Policlinico San Matteo
Molecular Virology Unit
Prof. Fausto Baldanti

Maurizio Zavattoni
Giuditta Comolli
Milena Furione
Stefania Paolucci
Antonella Sarasini
Elena Percivalle
Maurizio Parea
Giulia Campanini

Luca Dossena
Ermanna Cabano
Laura Bellotti
Claudia Lucchelli
Gabriella Garbagnoli
Viviana Landini
Simona Pezzaia
Giuliana Casali
Gabriella Frisco

Francesca Rovida
Irene Cassaniti
Federica Giardina
Federica Novazzi
Guy A.

